

日本大学生物資源科学部生命科学研究所
平成26年度発表論文一覧

1. Isobe K., Higo M., Kondo T., Sato N., Takeyama S. and Torigoe Y. Effect of winter crop species on arbuscular mycorrhizal fungal colonization and subsequent soybean yields. *Plant Production Science*, 17(3): 260-267, 2014.
2. Sakuma C., Furihara K., Nishio T., Miyakawa T., Tanokura M. and Tashiro M. Analysis of weak affinity of β -D-fructofuranosyl-(2 \rightarrow 1)-2-acetamido-2-deoxy- α -D-glucopyranoside for yeast β -fructofuranosidase using NMR spectroscopy. *Journal of Carbohydrate Chemistry*, 33: 498-505, 2014.
3. Hirano T., Sugiyama K., Sakaki Y., Hakamata W., Park S.-Y. and Nishio T. Structure-based analysis of domain function of chitin oligosaccharide deacetylase from *Vibrio parahaemolyticus*. *FEBS Letters*, 589: 145-151, 2015.
4. Minamoto T., Takahashi N., Kitahara S., Shinozaki Y., Hirano T., Hakamata W., Nishio T. Saccharification of β -chitin from squid pen by a fermentation method using recombinant chitinase-secreting *Escherichia coli*. *Applied Biochemistry and Biotechnology*, 175(8): 3788-3799, 2015.
5. Hakamata W., Tamura S., Hirano T., Nishio T. Multicolor imaging of endoplasmic reticulum-located esterase as a prodrug activation enzyme. *ACS Medicinal Chemistry Letters*, 5: 321-325, 2014.
6. Hakamata W., Miura K., Hirano T., Nishio T. Identification of a novel glycan processing enzyme with *exo*-acting β -allosidase activity in the Golgi apparatus using a new platform for the synthesis of fluorescent substrates. *Bioorganic Medicinal Chemistry*, 23: 73-79, 2015.
7. Okura K., Tamura R., Shigehara K., Masai E., Nakamura M., Otsuka Y., Katayama Y., Nakao Y. Synthesis of Polysubstituted Benzenes from 2-Pyrone-4,6-dicarboxylic Acid. *Chemistry Letters*, 43: 1349-1351, 2014.
8. Tamura M., Tsuji Y., Kusunose T., Okazawa A., Kamimura N., Mori T., Nakabayashi R., Hishiyama S., Fukuhara Y., Hara H., Sato-Izawa K., Muranaka T., Saito K., Katayama Y., Fukuda M., Masai E., Kajita S. Successful expression of a novel bacterial gene for pinoresinol reductase and its effect on lignan biosynthesis in transgenic *Arabidopsis thaliana*. *Appl Microbiol Biotechnol*, 98(19): 8165-8177, 2014.
9. Takahashi K., Kamimura N., Hishiyama S., Hara H., Kasai D., Katayama Y., Fukuda M., Kajita S., Masai E. Characterization of the catabolic pathway for a phenylcoumaran-type lignin-derived biaryl in *Sphingobium* sp. strain SYK-6. *Biodegradation*, 25(5): 735-745, 2014.

10. Yoshikata T., Suzuki K., Kamimura N., Namiki M., Hishiyama S., Araki T., Kasai D., Otsuka Y., Nakamura M., Fukuda M., Katayama Y., Masai E. Three-component *O*-demethylase system essential for catabolism of a lignin-derived biphenyl compound in *Sphingobium* sp. strain SYK-6. *Appl. Environ. Microbiol.*, 80(23): 7142-7153, 2014.
11. Endo H., Yamaguchi M., Tamura T., Nakano Y., Nishikubo N., Yoneda A., Kato K., Kubo M., Kajita S., Katayama Y., Ohtani M., Demura T. Multiple classes of transcription factors regulate the expression of *VASCULAR-RELATED NAC-DOMAIN7*, a master switch of xylem vessel differentiation. *Plant Cell Physiol.*, 56(2): 242-254, 2015.
12. Itonaga K. Contamination and community support in the aftermath of the Fukushima disaster. *Bulletin of the Atomic Scientists*, 0: 1-8, 2014.
13. Hirayama S., Shimizu M., Tsuchiya N., Furukawa S., Watanabe D., Shimoi H., Takagi H., Ogihara H., Morinaga Y. Awa1p on the cell surface of *sake* yeast inhibits biofilm formation and the co-aggregation between *sake* yeasts and *Lactobacillus plantarum* ML11-11. *Journal of Bioscience and Bioengineering*, 119(5): 532-537, 2015.
14. Furukawa S., Isomae R., Tsuchiya N., Hirayama S., Yamagishi A., Kobayashi M., Suzuki C., Ogihara H., Morinaga Y. Screening of lactic acid bacteria that can form mixed-species biofilm with *Saccharomyces cerevisiae*. *Bioscience, Biotechnology and Biochemistry*, 79(4): 681-686, 2015.
15. Watsuji T., Takano H., Yamabe T., Tamazawa S., Ikemura H., Ohishi T., Matsuda T., Shiratori-Takano H., Beppu T., Ueda K. Analysis of the tryptophanase expression in *Symbiobacterium thermophilum* in a coculture with *Geobacillus stearothermophilus*. *Appl Microbiol Biotechnol*, 98: 10177-10186, 2014.
16. Takano H., Hagiwara K., Ueda K. Fundamental role of cobalamin biosynthesis in the developmental growth of *Streptomyces coelicolor* A3 (2). *Appl Microbiol Biotechnol*, 99: 2329-2337, 2015.
17. Takano H., Agari Y., Hagiwara K., Watanabe R., Yamazaki R., Beppu T., Shinkai A., Ueda K. LdrP, a cAMP receptor protein/FNR family transcriptional regulator, serves as a positive regulator for the light-inducible gene cluster in the megaplasmid of *Thermus thermophilus*. *Microbiology*, 160: 2650-2660, 2014.
18. Shiratori-Takano H., Akita K., Yamada K., Itoh T., Sugihara T., Beppu T., Ueda K. Description of *Symbiobacterium ostreiconchae* sp. nov., *Symbiobacterium turbinis* sp. nov. and *Symbiobacterium terraclitae* sp. nov., isolated from shellfish, emended description of the genus *Symbiobacterium* and proposal of *Symbiobacteriaceae* fam. nov. *International Journal of Systematic and Evolutionary Microbiology*, 64: 3375-3383, 2014.

19. Wakabayashi T., Joseph B., Yasumoto S., Akashi T., Aoki T., Harada K., Muranaka S., Bamba T., Fukusaki E., Takeuchi Y., Yoneyama K., Muranaka T., Sugimoto Y., Okazawa A. Planteose as a storage carbohydrate required for early stage of germination of *Orobanche minor* and its metabolism as a possible target for selective control. *Journal of Experimental Botany*, 66(11): 3085-3097, 2015.
20. Sawai S., Ohyama K., Yasumoto S., Seki H., Sakuma T., Yamamoto T., Takebayashi Y., Kojima M., Sakakibara H., Aoki T., Muranaka T., Saito K., Umemoto N. Sterol side chain reductase 2 is a key enzyme in the biosynthesis of cholesterol, the common precursor of toxic steroidal glycoalkaloids in potato. *Plant Cell*, 26(9): 3763–3774, 2014.
21. Sawada Y., Aoki T. Metabolomics. In *The Lotus japonicus Genome* (Tabata S. and Stougaard J. eds.), Springer, 171-182, 2014.
22. Takihara H., Matsuura C., Ogihara J., Iwabuchi N., Sunairi M. *Rhodococcus rhodochrous* ATCC12674 becomes alkane-tolerant upon GroEL2 overexpression and survives in the *n*-octane phase in two phase culture. *Microbes and Environments*, 29(4): 431-433, 2014.
23. Takihara H., Ogihara J., Yoshida T., Okuda S., Nakajima M., Iwabuchi N., Sunairi M. Enhanced translocation and growth of *Rhodococcus erythropolis* PR4 in the alkane phase of aqueous-alkane two phase cultures were mediated by GroEL2 overexpression. *Microbes and Environments*, 29(4): 346-352, 2014.
24. Ogura T., Ogihara J., Sunairi M., Takeishi H., Aizawa T., Olivos-Trujillo MR., Maureira-Butler IJ., Salvo-Garrido HE. Proteomic characterization of seeds from yellow lupin (*Lupinus luteus* L.). *Proteomics*, 14(12): 1543-1546, 2014.
25. Wada N., Sunairi M., Anzai H., Iwata R., Yamane A., Nakajima M. Glycolytic activities in the larval digestive tract of *Trypoxylus dichotomus* (Coleoptera: Scarabaeidae). *Insects*, 5: 351-363, 2014.
26. Urai M., Aizawa T., Nakajima M., Sunairi M. An anionic polymer incorporating low amounts of hydrophobic residues is a multifunctional surfactant. Part 1: Emulsifying, thickening, moisture-absorption and moisture-retention abilities of a fatty acid-containing anionic polysaccharide. *Advances in Chemical Engineering and Science*, 5: 173-180, 2015.
27. Urai M., Aizawa T., Nakajima M., Sunairi M. An anionic polymer incorporating low amounts of hydrophobic residues is a multifunctional surfactant. Part 2: Emulsification, moisture absorption, and moisture retention of alkyl esterified poly- γ -glutamic acid. *Advances in Chemical Engineering and Science*, 5: 181-191, 2015.
28. 平野展孝. 駕籠に乗る人, 担ぐ人, そのまた草鞋を作る人. *生物工学会誌*. 第 92 卷 第 6 号: 305, 2014. 平野 1